

# Geometry

Anson Li

## Contents

<b>1</b>	<b>Manifolds</b>	<b>1</b>
1.1	Introduction	1
1.2	Structure of a manifold	2
1.3	Morphisms between manifolds	3
1.4	Tangent spaces	4
1.5	Velocities	5
1.6	Splitting	5
1.7	Submanifolds	6
1.8	Notes and References	8
<b>2</b>	<b>Vector Bundles</b>	<b>9</b>
2.1	Vector Bundles	9
2.2	Tangent Bundle	11
2.3	Notes and References	12
<b>3</b>	<b>Coordinates</b>	<b>14</b>
3.1	Introduction	14
3.2	Exterior Derivative	14
3.3	Derivations	14
3.4	Boundary	15

## 1. Manifolds

### 1.1. Introduction

We serve to give the reader the shortest introduction to manifold theory. This and the subsequent two chapters are loosely based on [Lan12; Lee13], and the symbols  $E, F$  will always denote Banach spaces, and all Banach spaces are assumed to be over  $\mathbb{R}$ . We sometimes say  $E$  (resp.  $F$ ) is a space for brevity, and

- $\mathcal{L}(E, F)$  = linear maps between  $E$  and  $F$ ,
- $L(E, F)$  = toplinear (continuous and linear) maps between  $E$  and  $F$ ,
- $\text{Topliso}(E, F)$  = toplinear isomorphisms between  $E$  and  $F$ ,
- $\text{Laut}(E)$  = toplinear automorphisms on  $E$ , which form a strongly open subset of  $L(E, E)$ .

We will be working in the category of  $C^p$  Banach spaces — where  $p \geq 0$ . The morphisms in the category of  $\text{Ban}_{\mathbb{R}}$  are called  $C^p$  morphisms, which are  $p$ -times continuously differentiable functions.

#### Definition 1.1 Morphisms between open subsets of Banach spaces

Let  $E$  and  $F$  be Banach spaces, and  $U \subseteq E$ ,  $V \subseteq F$  be open subsets. A mapping  $f : U \rightarrow F$  is of class  $C^p$  if  $f \in C(E, F)$  and Eq. (1) holds.

$$D^{(i)}f : U \rightarrow L^i(E, F) \quad \text{exists and is continuous for } i = p \quad (1)$$

$C^p(E, F)$  denotes the vector space of  $C^p$  mappings between  $E$  and  $F$ . Sometimes, we restrict our attention to *open subsets* of  $E$  and  $F$ , in this case:  $f \in C^p(U, V)$  if  $f \in C(U, V)$  and Eq. (2) holds.

$$D^{(i)}f : U \rightarrow L^i(E, F) \quad \text{exists and is continuous for } i = p \quad (2)$$

We sometimes write  $C^p$  for  $C^p(E, F)$  when it is clear. A  $C^p$  isomorphism is a bijective  $C^p$  morphism whose inverse is also a morphism.

#### Remark 1.2 Implicit assumption

In Eq. (2) we assumed that  $f(U) \subseteq V$ . This is a non-trivial part of the definition of  $C^p$  morphisms between  $E$  and  $F$ , we will come back to this in Def. 1.13.

Let  $f_1$  and  $f_2$  be mappings, and  $X$  a non-empty set.

- We say they are *composable* if either one of  $f_2 \circ f_1$  or  $f_1 \circ f_2$  makes sense.
- We also write  $f_2 f_1$  to refer to  $f_2 \circ f_1$  if there is no ambiguity.
- If  $U \subseteq X$  and  $V \subseteq Y$ , and  $f : U \rightarrow V$  is a bijection — meaning  $f(U) = V$  and  $f$  is injective, we say  $f$  is a bijection between  $U$  and  $V$ .
- With regards to inverse image notation, we allow ourselves to write

$$f_2^{-1} \circ f_1^{-1} \quad \text{is the same as} \quad f_2^{-1} f_1^{-1}$$

and inversion is never left associative.

$$f_2 f_1^{-1} = f_2 \circ f_1^{-1} \neq (f_2 \circ f_1)^{-1}$$

Composable  $C^p$  mappings are functors in the category of open subsets between Banach spaces. Few basic facts about  $C^p$  morphisms:

- If  $f$  is a toplinear mapping between  $E$  and  $F$ , then  $f \in C^p(E, F)$  for all  $p \geq 0$ .
- If  $f$  is a bijective toplinear mapping, then it is a  $C^p$  isomorphism for all  $p \geq 0$ .
- However, a bijective  $C^p$  morphism need not be a  $C^p$  isomorphism.

## 1.2. Structure of a manifold

It is fruitful to *construct* the manifold rather than *define* it. We also insist on working with open sets of Banach spaces instead coordinate functions as our primary data.

### Definition 1.3 Chart

Let  $X$  be a non-empty set. A *chart on  $X$  modelled on a Banach space  $E$*  is a tuple  $(U, \varphi)$ , such that  $U \subseteq X$ ,  $\varphi(U) = \hat{U}$  is an *open* subset of  $E$ , and  $\varphi$  is a bijection onto  $\hat{U}$ .

### Definition 1.4 Compatibility

Let  $(U, \varphi)$  and  $(V, \psi)$  be charts on  $X$  modelled on  $E$ , they are called  $C^p$  compatible (for  $p \geq 0$ ) if  $U \cap V = \emptyset$ , or both of the following hold

- $\varphi(U \cap V)$  and  $\psi(U \cap V)$  are *both* open subsets of  $E$ , and
- the *transition map*  $\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V)$  is a  $C^p$  isomorphism between open subsets of  $E$ .

### Definition 1.5 Atlas

Let  $X$  be a non-empty set and  $p \geq 0$ . A  $C^p$  *atlas on  $X$  modelled on  $E$*  is a pairwise  $C^p$  compatible collection of charts  $\{(U_\alpha, \varphi_\alpha)\}$  whose union over the domains cover  $X$ .

We will assume hereinafter that atlases are of class  $C^p$  for  $p \geq 0$ . Let  $X$  be a non-empty set, equipped with an atlas  $\{(U_\alpha, \varphi_\alpha)\}$  modelled on a space  $E$ . Suppose  $\alpha$ , and  $\beta$  both index the atlas.

- We write  $\hat{U}_\alpha$  to refer to  $\varphi_\alpha(U_\alpha)$ , and
- $\hat{p} = \varphi_\alpha(p)$  for  $p \in U_\alpha$  when it is clear which chart we are using.
- $U_{\alpha\beta} = U_\alpha \cap U_\beta$ , and if  $U_{\alpha\beta} \neq \emptyset$ : the *transition map from  $\alpha$  to  $\beta$*  is defined in Eq. (3).

$$\varphi_{\alpha\beta} \triangleq \varphi_\beta|_{U_{\alpha\beta}} \circ (\varphi_\alpha|_{U_{\alpha\beta}})^{-1} : \varphi_\alpha(U_{\alpha\beta}) \rightarrow \varphi_\beta(U_{\alpha\beta}) \quad (3)$$

- We often suppress the restrictions of the two charts in the composition, and Eq. (3) reads

$$\varphi_{\alpha\beta} = \varphi_\beta \circ \varphi_\alpha^{-1} = \varphi_\beta \varphi_\alpha^{-1} \quad (4)$$

### Remark 1.6 Omissions of $C^p$

We might refer to two charts as *compatible* or *smoothly compatible*, implying they are  $C^p$  compatible. This comes from the perspective that, in the context of  $C^p$  manifolds, any smoothness exceeding  $C^p$  is deemed sufficiently smooth for our purposes. We also say  $C^p$  for  $C^p$  where  $p \geq 0$ .

### Definition 1.7 Structure determined by an atlas

Let  $\mathcal{A}$  be an atlas on  $X$ , the maximal atlas containing  $\mathcal{A}$  is called the  $C^p$  *structure determined by  $\mathcal{A}$* .

### Definition 1.8 Manifold

A  $C^p$  *manifold modelled on  $E$*  is a non-empty set  $X$  with a  $C^p$  structure modelled on  $E$ . We refer to  $E$  as the *model space of  $X$* .

### Proposition 1.9 $E$ is a manifold

The identity  $\text{id}_E$  defines an atlas on  $E$ , which determines a  $C^p$  structure called the *standard structure of  $E$*  for  $p \geq 0$ . We call  $(E, \text{id}_E)$  the *standard chart on  $E$* .

### Proposition 1.10 Topology is unique on a manifold

Let  $X$  be a  $C^p$  manifold modelled on  $E$ , it induces a unique topology such that the domain for each chart in its smooth structure is open, and each chart is a homeomorphism onto its range in the subspace topology.

*Proof.* We offer a sketch of the proof. Fix a chart  $(U, \varphi)$ , it is clear that  $U$  has to be in the topology of  $X$ , and because  $\varphi : U \rightarrow \hat{U}$  is required to be a homeomorphism, we duplicate all the open sets in  $\hat{U}$  by using the inverse image through  $\varphi$ . The collection of all such inverse images form a sub-basis, thus defines a unique topology as is well known.

There is an alternate way constructing the above topology. The existence of a coarsest topology on a chart domain  $U$  such that if  $(V, \varphi)$  is another chart whose domain  $V \cap U \neq \emptyset$ , then  $\varphi|_{U \cap V}$  is a homeomorphism onto its range. Stitching the weak topologies together, we obtain an ambient topology on  $X$  [Lan12]. ■

**Remark 1.11 Omission of model space**

For any of the objects we have defined in this section, that depend upon a model space or a morphism class (i.e  $C^p$ ), we will say 'X is a manifold', rather than X is a manifold of class  $C^p$  modelled over E when it is convenient to do so. If the model space E is infinite (resp. finite) dimensional, we say X is infinite (resp. finite) dimensional. And a reminder:  $C^p$  should always be interpreted with  $p \geq 0$ .

**Proposition 1.12 Open subsets of manifolds**

Let  $U$  be an open subset of a manifold  $X$ , then  $U$  is a manifold whose structure is determined by the atlas  $\mathcal{A}$  in Eq. (5).

$$\mathcal{A} = \{(V, \varphi) \text{ in the structure of } X, \text{ where } V \subseteq U\} \quad (5)$$

*Proof.* The structure of  $X$  includes all possible restrictions to open sets; hence  $\mathcal{A}$  in Eq. (5) is an atlas, and a unique structure by Def. 1.7. ■

### 1.3. Morphisms between manifolds

**Definition 1.13 Morphisms between manifolds**

A mapping  $f : X \rightarrow Y$  between manifolds is a *morphism* (a  $C^p$  morphism to be precise) if for every  $p \in X$ , there exist charts  $(U, \varphi) \in X$  and  $(V, \psi) \in Y$  such that 1) the image  $f(U)$  is contained in a the chart domain  $V$ , and 2)

$$f_{U,V} \stackrel{\Delta}{=} \psi \circ f \circ \varphi^{-1} \in C^p(\hat{U}, \hat{V}) \quad \text{in the sense of Def. 1.1.} \quad (6)$$

The map  $f_{U,V}$  as defined in Eq. (6) is called the *coordinate representation* of  $f$  with respect to the charts  $(U, \varphi), (V, \psi)$ .

**Remark 1.14 Identifying  $X$  with its structure**

If  $(U, \varphi)$  is a chart in the structure of  $X$ , we will simply say  $(U, \varphi)$  is in  $X$ .

**Remark 1.15 Identifying charts with their domains**

The scenario in Eq. (6) occurs so often that we decide to simply write

$$f_{U,V} = \psi f \varphi^{-1} \quad (7)$$

to mean there exists charts  $(U, \varphi), (V, \psi)$  in the structure of  $X, Y$  with

$$f(U) \subseteq V \quad (8)$$

Consistent with the notation of putting hats on objects

borrowed or pulled back from the model spaces, we write  $\hat{f} = f_{U,V}$ . Equation (9) gives an example of this.

$$\hat{f}(\hat{p}) = f_{U,V}(\hat{p}) = f_{U,V}(\varphi(p)) \quad (9)$$

for any morphism  $f \in \text{Mor}(X, Y)$ , and charts that satisfy Eq. (8). We refer to the map in Eq. (9) as a *coordinate representation of  $f$  about  $p$* , with the inference that  $p \in (U, \varphi)$ .

Definition 1.13 may leave one unsatisfied. Why do we require the image  $f(U)$  be contained in another chart domain in  $Y$ ? There are two reasons.

**Continuity** Suppose  $f$  is a map between  $E$  and  $F$ , and the restriction of  $f$  onto a family of open subsets  $U_\alpha \subseteq E$  is  $C^p$  for  $p \geq 0$ . If  $\{U_\alpha\}$  is an open cover for  $E$ , then  $f$  is continuous. Proposition 1.16 shows this equally holds for manifolds.

**Implicit assumption** The definition of smoothness between open subsets of Banach spaces (see Def. 1.1) is a purely local one. And let us recall: every chart domain  $U$  in a manifold  $X$  corresponds to an open subset  $\hat{U} \subseteq E$  in the model space, and see Rmk 1.2 as well. Hence, **the necessity that the image  $f(U)$  is contained in a single chart domain of  $Y$  is a relic of the original definition**. The astute reader will also see that the openness requirement of  $\psi(U \cap V)$ , and  $\varphi(U \cap V)$  in Def. 1.4 is completely natural as well, since  $C^p$  morphisms are defined between open subsets of Banach spaces.

**Proposition 1.16 Properties of morphisms between manifolds**

Every  $C^p$  morphism between manifolds is a continuous map, and the composition of  $C^p$  morphisms is again a morphism.

*Proof.* The first claim is proven if we show  $f$  is locally continuous. Using Equation (6), since  $p$  is arbitrary, choose any neighbourhood  $W$  of  $f(p)$ , by shrinking this neighbourhood, it suffices to assume it is a subset of the chart domain  $V$ . The charts on  $X$  and  $Y$  are homeomorphisms, and unwinding the formula shows that  $f|_U = \psi^{-1} f_{U,V} \varphi$ , so that

$$U \cap f^{-1}(W) = (f|_U)^{-1}(W) \quad \text{is open in } X.$$

To prove the second, let  $X_3$  be manifolds modelled over  $E_3$ , and  $f_1, f_2$  is smooth between  $X_i$  such that  $f_2 \circ f_1$  makes sense. Since  $f_1$  is smooth, one finds a pair of charts  $(U_i, \varphi_i) \in X_i$  for  $i = 1, 2$  about each  $p \in X_1$  such that  $(f_1)_{U_1, U_2}$  is a morphism.

$f_2(f_1(p))$  induces another pair of charts  $(V_i, \psi_i) \in X_i$  for  $i = 2, 3$ . Since  $f_2$  is smooth, it is continuous.  $f_1^{-1} \circ f_2^{-1}(V_3)$  is open in  $X_1$ , and we can shrink all of our charts so that

$f_2 f_1(U_1)$  is contained in  $V_3$ . Finally, because  $C^p$  morphisms between open subsets of Banach spaces is closed under composition,  $f_{U_1 \cap f_1^{-1} f_2^{-1}(V_3), V_3}$  is smooth. ■

### Remark 1.17 Morphisms between $C^k, C^p$ manifolds

Let  $X$  be a  $C^k$ -manifold, and  $Y$  a  $C^p$  manifold, where  $k, p \geq 0$ . A morphism between  $X$  and  $Y$  is a map  $f : X \rightarrow Y$  such that each point  $p \in X$  admits a coordinate representation

$$f_{U,V} \in C^{\min(p,k)}(\hat{U}, \hat{V}) \quad (10)$$

If  $\min(p, k) \geq 1$ , then we define its differential as in Def. 1.21 by treating both  $X$  and  $Y$  as  $C^{\min(k,p)}$  manifolds.

## 1.4. Tangent spaces

**In this section, all manifolds will be of class  $C^p$  for  $p \geq 1$ .** The next question that we will address is taking derivatives of smooth maps between manifolds. There is no reason to demand  $C^p$  smoothness between maps, or even a  $C^p$  category of manifolds if we cannot borrow something more other than the morphisms on open sets.

Suppose  $U$  is an open subset of  $E$  and  $f : U \rightarrow Y$  is  $C^p$ . The derivative  $Df(x)$  is a linear map  $E \rightarrow F$ , not from  $U$  to  $F$  ( $U$  might not even be a vector space). This suggests the 'derivative' of a morphism  $F : X \rightarrow Y$  between manifolds can in some sense be interpreted as the *ordinary derivative* of its coordinate representation  $DF_{U,V}(\hat{p})$ , adhering to our principle of using open sets.

But there is a problem with this 'derivative': it gives different values for different charts. With infinitely many charts in  $X$  and  $Y$ , this definition becomes useless. To see this, let  $X$  be a manifold modelled on  $E$  and  $p \in X$ . If  $f : X \rightarrow Y$  is a morphism, and  $(U_1, \varphi_1), (U_2, \varphi_2)$  are charts defined about  $p$  such that the representations  $f_{U_1, V}$  and  $f_{U_2, V}$  are morphisms. Writing  $p_i = \varphi_i(p)$ ,  $U_{12} = U_1 \cap U_2$  and

$$\varphi_{12} = \varphi_2 \varphi_1^{-1} : \varphi_1(U_{12}) \rightarrow \varphi_2(U_{12}) \quad (11)$$

(because the map in Eq. (11) goes from the domain  $U_1$  to  $U_2$ ), a simple computation yields Eq. (12).

$$\begin{aligned} Df_{U_1, V}(p_1)(v) &= D(\psi f \varphi_2^{-1} \varphi_2 \varphi_1^{-1})(p_1)(v) \\ &= Df_{U_2, V}(p_2)(D\varphi_{12}(p_1)(v)) \\ &= Df_{U_2, V}(p_2) \circ D\varphi_{12}(p_1) \cdot (v) \end{aligned} \quad (12)$$

where  $\cdot(v)$  denotes the evaluation at  $v \in E$ , and is assumed to be left associative over composition. The computation

in Eq. (12) suggests that interpreting the derivative by pre-conjugation is dependent on the chart being used to interpret the derivative. In fact,  $D\varphi_{12}(p_1)$  can be replaced with any toplinear isomorphism on  $E$  (relabel  $\varphi_2 = A\varphi_1$  where  $A$  is any linear automorphism on  $E$ ), so the right hand side of Eq. (12) can be interpreted as  $Df_{U_2, V}(p_2)(w)$  where  $w$  is any vector in  $E$ .

### Definition 1.18 Concrete tangent vector

Suppose  $k \geq 1$ ,  $X$  a  $C^k$ -manifold on  $E$ , and  $p \in X$ . If  $(U, \varphi)$  is any chart containing  $p$ , for each  $v \in E$  we call  $(U, \varphi, p, v)$  a *concrete tangent vector at  $p$*  that is *interpreted* with respect to the chart  $(U, \varphi)$ . The disjoint union of concrete tangent vectors, as shown in Eq. (13)

$$T_{(U, \varphi, p)} X = \bigcup_{v \in E} \{(U, \varphi, p, v)\} \cong E, \quad (13)$$

is called the *concrete tangent space at  $p$*  interpreted with respect to  $(U, \varphi)$ ; and it inherits a TVS structure from  $E$ .

Fix a point  $p$  in a manifold  $X$ . Suppose  $(U_i, \varphi_i)$  are charts containing  $p$ , from Eq. (12) there exists a natural (toplinear) isomorphism between the concrete tangent spaces, namely

$$(U_1, \varphi_1, p, v_1) \sim (U_2, \varphi_2, p, v_2) \quad \text{iff} \quad v_2 = D\varphi_{12}(p_1)(v_1), \quad (14)$$

where  $p_i = \varphi_i(p)$ . The right member of Eq. (14) is the derivative of a transition map — which is a toplinear automorphism on  $E$ . Hence  $D\varphi_{12}(p_1)$  defines a toplinear isomorphism between  $T_{(U_1, \varphi_1, p)} X$  and  $T_{(U_2, \varphi_2, p)} X$ . With this, we define the primary object of our study.

### Definition 1.19 Tangent vector

A *tangent vector* (or an *abstract tangent vector*) at  $p$  is defined as an equivalence class of concrete tangent vectors at  $p$ , under the relation in Eq. (14).

### Definition 1.20 Tangent space

The *tangent space at  $p$* , denoted by  $T_p X$  is the set of all tangent vectors at  $p$ . It is toplinearly isomorphic to the model space  $E$ .

### Definition 1.21 Differential of a morphism

Let  $X$  and  $Y$  be modelled on the spaces  $E$  and  $F$ . If  $f$  be a morphism between  $X$  and  $Y$ , and fix  $p \in X$ . We define a linear map, called the *differential of  $f$  at  $p$*  shown in Eq. (15).

$$df(p) : T_p X \rightarrow T_{f(p)} Y \quad (15)$$

Whose action on tangent vectors is characterized by

- if  $(U, \varphi)$  and  $(V, \psi)$  are any pair of charts that satisfy the morphism condition in Eq. (6) about  $p$ , and suppose
- $v \in T_p M$  is represented by  $(U, \varphi, p, \hat{v})$
- then  $df(p)(v) \in T_{f(p)} Y$  is represented by  $(V, \psi, f(p), Df_{U,V}(\hat{p})(\hat{v}))$

Alternatively, the diagram shown in fig. 1 commutes. We also write  $df_p = df(p)$ .

$$\begin{array}{ccc} T_p X & \xrightarrow{\quad} & T_{(U,\varphi,p)} X \\ \downarrow df(p) & & \downarrow Df_{U,V}(\hat{p}) \\ T_{f(p)} Y & \xrightarrow{\quad} & T_{(V,\psi,f(p))} Y \end{array}$$

Figure 1: Differential of a morphism

## 1.5. Velocities

In the previous section, we motivated the definition of  $T_p X$  using the computation of the derivative of a morphism from  $X$ . Dually, the tangent space allows us compute the derivatives of morphisms into  $X$  in a coordinate independent manner.

### Definition 1.22 Curve

Let  $J_\varepsilon = (-\varepsilon, +\varepsilon)$  be an open interval in  $\mathbb{R}$  containing the origin. Proposition 1.12 tells us  $J_\varepsilon$  is a manifold. A morphism  $\gamma: J_\varepsilon \rightarrow X$  is called a *curve in  $X$* , and  $\gamma(0)$  is called the *starting point of  $\gamma$* .

### Remark 1.23 Omission of chart in concrete representation

If  $p$  is a point on a manifold  $X$ , and  $v \in T_p X$  is represented by  $(U, \varphi, p, \hat{v})$ , we write

$$(U, \hat{v}) = (\hat{p}, \hat{v}) = \hat{v} = (U, \varphi, p, \hat{v}) \quad (16)$$

### Remark 1.24 Standard representation of tangent vectors

If  $X$  is an open subset of  $E$ , and  $p \in X$ , we identify a tangent vector  $v \in T_p X$  by its *standard representation*. Instead of using a  $\hat{v}$ , we use  $\bar{v}$ .

$$(X, \text{id}_X, p, \bar{v}) = (X, \bar{v}) = (X, \hat{v}) \quad \text{is a representation of } v \in T_p X \quad (17)$$

### Definition 1.25 Velocity of a curve

Let  $\gamma$  be a curve in  $X$  and  $t \in J_\varepsilon$ . We denote the *velocity* of a curve  $\gamma$  at  $t = t_0$  by  $\gamma'(t_0)$ ; which is defined in Eq. (18).

$$\gamma'(t_0) = [D\gamma_{J_\varepsilon, V}(t_0)(\bar{1})] \quad (18)$$

where  $(J_\varepsilon, \text{id}_{J_\varepsilon}, t_0, \bar{1})$  is a concrete tangent vector within  $T_{t_0} J_\varepsilon$ .

Equation (18) might seem arbitrary at first, but we must emphasize that this is the most natural interpretation of a velocity, encodes much of the geometric information of a tangent vector. We will revisit this topic when we discuss exterior differentiation.

### Proposition 1.26 Tangent vectors are velocities

Let  $p$  be a point on a manifold  $X$ . For every tangent vector  $v \in T_p X$ , there exists a curve starting at  $p$  whose velocity is  $v$ .

*Proof.* Find a chart  $(U)$  in  $X$  where  $\hat{p} = 0$ . Such a chart exists, because translations and dilations are  $C^p$  isomorphisms. If the tangent vector  $v$  has interpretation  $\hat{v}$  in  $U$ , there exists  $\varepsilon > 0$  so small that the range of  $\hat{\gamma}$ , as defined Eq. (19), lies in  $\hat{U}$

$$\hat{\gamma}: J_\varepsilon \rightarrow \hat{U} \quad \text{where} \quad \hat{\gamma}(t) = \int_0^t \hat{v} dt \quad (19)$$

$\hat{\gamma}$  is a curve in  $\hat{U}$  starting at  $\hat{p}$  with velocity  $\hat{v}$ . Defining  $\gamma$  as the composition of  $\hat{\gamma}$  with the chart inverse finishes the proof. ■

## 1.6. Splitting

Recall: if  $W$  is a vector space and  $W_1, W_2$  are linear subspaces of  $V$ .  $W_2$  is the vector space complement of  $W_1$  (resp. with the indices reversed) if

$$W_1 + W_2 = W, \quad \text{and} \quad W_1 \cap W_2 = 0$$

We sometimes refer to the vector space complement of  $W_1$  as its *linear complement*.

### Definition 1.27 Splitting in $E$

A linear subspace  $E_1$  splits in  $E$  if both  $E_1$  and its vector space complement  $E_2$  are closed, and the addition map  $\theta: E_1 \times E_2 \rightarrow E$  given by

$$\theta(x, y) = x + y \quad \text{is a toplinear isomorphism.}$$

### Definition 1.28 Splitting in $L(E, F)$

A continuous, injective linear map  $\lambda \in L(E, F)$  splits iff

its range splits in  $F$ .

Every finite dimensional or finite codimensional linear subspace of  $E$  splits. And if  $E$  itself is finite dimensional, then every linear subspace of  $E$  splits. An alternative definition of Def. 1.28 is as follows: a map  $\lambda \in L(E, F)$  splits iff there exists a toplinear isomorphism  $\theta : F \rightarrow F_1 \times F_2$  such that  $\lambda$  composed with  $\alpha$  induces a toplinear isomorphism from  $E$  onto  $F_1 \times 0$  — which we identify with  $F_1$ .

If  $E$  and  $F$  are finite dimensional (so  $E = \mathbb{R}^n$  and  $F = \mathbb{R}^m$  respectively), Def. 1.28 refers to the familiar matrix canonical form in Eq. (20), and Defs. 1.29 and 1.30 can be seen as infinite-dimensional analogues of Eq. (20).

$$A_{\text{injective}} = \begin{bmatrix} \text{id}_{m \times m} \\ 0_{n-m \times m} \end{bmatrix} \quad A_{\text{surjective}} = [\text{id}_{n \times n} \quad 0_{n \times m-n}] \quad (20)$$

### Definition 1.29 Immersion

A morphism  $f \in \text{Mor}(X, Y)$  is an *immersion at  $p$*  if there exists a coordinate representation about  $f_{U,V}$  such that

$$Df_{U,V}(\hat{p}) \quad \text{is injective and splits.} \quad (21)$$

The morphism  $f$  is called an *immersion* if Eq. (21) holds at every  $p$ .

### Definition 1.30 Submersion

A morphism  $f \in \text{Mor}(X, Y)$  is a *submersion at  $p$*  if there exists a coordinate representation about  $f_{U,V}$  such that

$$Df_{U,V}(\hat{p}) \quad \text{is surjective and its kernel splits.} \quad (22)$$

The morphism  $f$  is called a *submersion* if Eq. (22) holds at every  $p$ .

### Definition 1.31 Embedding

A morphism  $f \in \text{Mor}(X, Y)$  is an *embedding* if it is an immersion and a homeomorphism onto its range.

### Definition 1.32 Toplinear subspace

Let  $E$  be a Banach space, a *toplinear subspace of  $E$*  is a closed linear subspace  $E_1$  which splits in  $E$ .

## 1.7. Submanifolds

Before we state the definition of a submanifold, it is important to recapitulate the construction of a manifold  $X$ .

- Given a non-empty set  $X$  and an atlas modelled on a space  $E$ .
- The purpose of each chart in the atlas is to borrow open subsets  $\hat{U} \subseteq E$ . If we single out a single chart, **the construction is entirely topological**. It is of little importance *how* the individual chart domains  $U$  are mapped onto  $\hat{U}$ ,
- Each chart is in **bijection with its range**, which is an open subset of  $E$ , and
- the transition maps  $\varphi_{\alpha\beta} = \varphi_\beta \varphi_\alpha^{-1}$  are **morphisms between open subsets of  $E$** .

If  $(U, \varphi) \in X$  is a chart whose domain intersects  $S$ , the question then becomes: Is it possible to modify  $(U, \varphi)$  so that it becomes a chart modelled on  $E_1$ ? If we restrict  $\varphi$  onto  $U \cap S$ , its range is still an open subset of  $E$ . We can assume  $\varphi(U \cap S) \subseteq E$  is constant on the linear complement of  $E_1$ , that way  $\varphi|_{U \cap S}$  will be a bijection.

The range of the restricted chart is still a subset of  $E$ , and not  $E_1$ . An easy fix to this would be to require  $E_1$  to **split in  $E$**  (and shrinking  $U$  using a basis argument). Let  $\theta$  be a toplinear isomorphism between  $E$  and  $E_1 \times E_2$ , and we obtain Eq. (23).

$$\theta\varphi(S \cap U) = \hat{U}_1 \times a_2 \quad \text{where} \quad \hat{U}_1 \subseteq E_1 \text{ and } a_2 \in E_2. \quad (23)$$

Identifying  $\hat{U}$  with  $\theta(\hat{U})$ , and requiring  $U_1 \times a_2$  to be in  $\theta(\hat{U})$ , we arrive at the following definition.

### Definition 1.33 Submanifold

Let  $X$  be a manifold, and  $S$  a subset of  $X$ . We call  $S$  a *submanifold of  $X$*  if

- there exist split subspaces  $E_1, E_2$  of  $E$ ,
- for every  $p \in S$  is contained in  $(U, \varphi) \in X$  — a chart which also splits:

$$\varphi : U \rightarrow \hat{U} \cong \hat{U}_1 \times \hat{U}_2, \quad \text{where} \quad \hat{U}_i \subseteq E_i \quad \text{for } i = \underline{2}, \quad (24)$$

- and for a such chart, there exists  $a_2 \in \hat{U}_2$ , that makes  $U \cap S$  behave like a level set:

$$\varphi(U \cap S) = \hat{U}_1 \times a_2. \quad (25)$$

### Definition 1.34 Slice chart

We call a chart satisfying Eqs. (24) and (25) a *slice chart* of  $S$ ; and to simplify what follows, we write  $\varphi^i = \text{proj}_i \varphi$  for  $i = \underline{2}$  for any slice chart  $(U)$ .

Given that  $\text{proj}_i$  is a morphism between open subsets of Banach spaces,  $\varphi^i$  is again a morphism. In particular,  $\varphi^1$

is a bijection from  $U^s = U \cap S$  onto  $\hat{U}_1$ ; the latter being an open subset of  $E_1$ .

To show  $S$  is indeed a manifold it remains to show the collection of charts in Eq. (26) forms a  $C^p$  atlas modelled  $E_1$ , which we will prove in Proposition 1.35

$$\mathcal{A} = \{(U^s, \varphi^s) = (U^s, \varphi^1), (U, \varphi) \text{ is a slice chart of } S\}. \quad (26)$$

### Proposition 1.35 Structure of a submanifold

If  $S$  is a submanifold of  $X$ , Eq. (26) defines a  $C^p$  atlas over the space  $E_1$ . The manifold  $S$  has a topology that coincides with the subspace topology. Furthermore, the inclusion map  $\iota_S : S \rightarrow X$  is a morphism and an embedding.

*Proof.* Each of the charts in Eq. (26) is in bijection with an open subset of  $E_1$ . Let  $(U_\alpha^s, \varphi_\alpha^s)$  and  $(U_\beta^s, \varphi_\beta^s)$  be overlapping charts in  $\mathcal{A}$ . Using  $\theta$  as our toplinear isomorphism from  $E$  onto  $E_1 \times E_2$  as usual.

- By Eq. (24),  $(U_\alpha^s, \varphi_\alpha^s)$  is induced by a chart  $(U_\alpha, \varphi_\alpha) \in X$ ,  $\varphi_\alpha : U_\alpha \rightarrow \hat{U}_\alpha \subseteq E$  which splits into  $\theta(\hat{U}_\alpha) = \hat{U}_\alpha^s \times \hat{U}_{2,\alpha}$  such that  $\hat{U}_\alpha^s \subseteq E_1$  and  $\hat{U}_{2,\alpha} \subseteq E_2$ . Similarly for  $\beta$  as well.
- There exists elements  $a_2 \in \hat{U}_{2,\alpha}$ , (resp.  $b_2 \in \hat{U}_{2,\beta}$ ) where  $\theta\varphi_\alpha(U_\alpha^s) = \hat{U}_\alpha^s \times a_2$  resp.  $\beta$ .

### Note 1.36

Let us define  $U_{\alpha\beta}^s = U_\alpha^s \cap U_\beta^s$ , we will show Lem. 1.37.

### Lemma 1.37

Both  $\varphi_\alpha^s(U_{\alpha\beta}^s)$  and  $\varphi_\beta^s(U_{\alpha\beta}^s)$  are open subsets of  $E_1$ .

*Proof of Lem. 1.37.* We can factor  $U_{\alpha\beta}^s = (U^s \cap U_\alpha) \cap U_{\alpha\beta}$ , and because  $\varphi_\alpha$  is a bijection, we have  $\varphi_\alpha^s(U_{\alpha\beta}^s) = \text{proj}_1 \theta(\varphi_\alpha(U^s \cap U_\alpha) \cap \varphi_\alpha(U_{\alpha\beta}))$ .

$\theta$  and  $\text{proj}_1$  are both open maps, and because  $W \triangleq \varphi_\alpha(U_{\alpha\beta})$  is open in  $E$ :  $\theta(\varphi_\alpha(U^s \cap U_\alpha) \cap W)$  splits into a subset of  $\hat{U}_\alpha^s \times a_2$ ,

$\text{proj}_1 \theta(\varphi_\alpha(U^s \cap U_\alpha) \cap W) = \text{proj}_1(\text{Open subset of } E_1 \times a_2)$  which is open in  $E_1$ . ■

The diagram in fig. 2 provides a summary.

$$\begin{array}{ccccccc} U_{\alpha\beta}^s & \xrightarrow{\varphi_\alpha} & \varphi_\alpha(U_{\alpha\beta}^s) & \xrightarrow{\theta} & \varphi_\alpha(U_{\alpha\beta}^s)_1 \times a_2 & \xrightarrow{\text{proj}_1} & \varphi_\alpha^s(U_{\alpha\beta}^s) \\ \downarrow \varphi_{\alpha\beta} & & \downarrow & & \downarrow \theta\varphi_{\alpha\beta}\theta^{-1} & & \downarrow \\ U_{\alpha\beta}^s & \xrightarrow{\varphi_\beta} & \varphi_\beta(U_{\alpha\beta}^s) & \xrightarrow{\theta} & \varphi_\beta(U_{\alpha\beta}^s)_1 \times b_2 & \xrightarrow{\text{proj}_1} & \varphi_\beta^s(U_{\alpha\beta}^s) \end{array}$$

Figure 2: Overlap of slice charts

Identifying  $a_2$  (resp.  $b_2$ ) with the constant function ( $p \mapsto a_2$ ) for  $p \in U_\alpha^s$ , we get Eq. (27).

$$\varphi_\alpha^s \times a_2 = \theta \circ \varphi_\alpha \quad \text{resp. } \beta \quad (27)$$

Suppressing the restrictions onto domains, the transition map is given by the composition of maps in Eq. (28).

$$\varphi_\beta^s \circ (\varphi_\alpha^s)^{-1} = \text{proj}_1 \theta \varphi_\beta \varphi_\alpha^{-1} \theta^{-1} \text{proj}_1^{-1} : \varphi_\alpha^s(U_{\alpha\beta}^s) \rightarrow \varphi_\beta^s(U_{\alpha\beta}^s) \quad (28)$$

which is clearly a bijection. It suffices to show Eq. (28) is a morphism between open subsets of  $E_1$ . Let  $a_2 : \varphi_\alpha^s(U_{\alpha\beta}^s) \rightarrow \hat{U}_{2,\alpha}$ , which is the constant function  $a_2$  and hence a morphism.

The product  $(\text{id}_{\varphi_\alpha^s(U_{\alpha\beta}^s)} \times a_2) = \text{proj}_1^{-1}$  is a morphism into  $\varphi_\alpha^s(U_{\alpha\beta}^s) \times \hat{U}_{2,\alpha}$ . The inverse of  $\theta$  is an open morphism, and the terms  $\varphi_\beta \varphi_\alpha^{-1}$  combine into the transition map  $\varphi_{\alpha\beta}$  in  $X$  (up to a restriction on an open set). Equation (28) then reads

$$\varphi_\beta^s \circ (\varphi_\alpha^s)^{-1} = \text{proj}_1 \theta \varphi_{\alpha\beta} \theta^{-1} (\text{id}_{\varphi_\alpha^s(U_{\alpha\beta}^s)} \times a_2) \quad (29)$$

which is a morphism between open subsets. Reversing the roles of  $\alpha, \beta$  shows that Eq. (28) is an isomorphism. Therefore the collection of charts in Eq. (26) forms an atlas of  $S$ .

Let us use  $\iota_S : S \rightarrow X$  to represent the inclusion map and consider a point  $p \in S$ . It is always possible to identify a slice chart  $(U, \varphi)$  within  $X$  that contains  $p = \iota_S(p)$  in its domain. By definition of the atlas on  $S$ , this induces a truncated chart  $(U^s, \varphi^s)$ .

Observing that  $\iota_S(U^s) = \iota_S(U \cap S)$  lies within  $(U, \varphi)$ , the morphism criteria in Eq. (6) is satisfied. Computing the coordinate representation of  $\iota_S$ , we obtain Eq. (30).

$$(\iota_S)_{U^s, U} = \varphi \iota_S(\varphi^s)^{-1} = \text{id}_{\hat{U}_1} \times a_2 \quad (30)$$

Equation (30) shows that the coordinate representation of  $\iota_S$  is a local isomorphism. Since the inclusion map is a bijection and continuous, and the coordinate representation of  $\iota_S^{-1}$  is simply the inverse Eq. (30); and  $\iota_S^{-1}$  is a morphism and therefore continuous. The manifold topology of  $S$  coincides with its subspace topology.

At last, the inclusion map  $\iota_S$  has coordinate representation Eq. (30). Computing its ordinary derivative we obtain Eq. (31).

$$D(\iota_S)_{U^s, U}(\hat{p}) : T_{(U^s, \varphi^s, p)} \longrightarrow T_{(U, \varphi, p)}, \quad (31)$$

and  $D(\iota_S)_{U^s, U}(\hat{p}) = \text{id}_{E_1} \times 0$ . Which is a toplinear morphism between concrete tangent spaces and has a simple representation of 'adding zeroes' (see Def. 1.28) at the end of a vector  $\hat{v} \in E_1$  — which is to say: **the differential of  $\iota_S$  is injective and splits in  $E$** . Therefore  $\iota_S$  is an embedding. ■

#### Remark 1.38 Pairs of slice charts

Proposition 1.35 shows every point  $p \in S$  is in the domain of a slice chart in  $S$ , and the domain of the chart in  $X$  that induces the slice chart — whose inclusion map satisfies Eqs. (30) and (31). If  $p$  is a point on a submanifold  $S$ , we refer to a *pair of slice charts* containing  $p$  as the pair  $(U^s, \varphi^s)$  and  $(U, \varphi)$  in the structure of  $S$  and  $X$ .

#### Definition 1.39 Exterior tangent space of $S$

The *exterior tangent space* of a point  $p \in S$  is the image of  $T_p S$  under  $d\iota_S(p)$ ,

$$T_p^{\text{ext}} S = d\iota_S(p)(T_p S), \quad (32)$$

which is a toplinear subspace of  $T_p X$ .

## 1.8. Notes and References

#### Remark 1.40 Not necessarily Hausdorff

The topology generated by Prop. 1.10 is not necessarily Hausdorff, nor second countable. So a manifold  $X$  may not admit partitions of unity, but for our current purposes we will work with this general definition. Because of the uniqueness of the topology, we sometimes refer to the topology as being part of the *structure* of the manifold.

Lee [Lee13, Lemma 1.35] requires the chart domains separate points, and this leads to a Hausdorff topology. For a discussion on inductive systems and final topologies, see [Conway\_2019].

## 2. Vector Bundles

### 2.1. Vector Bundles

Let  $X$  be a class  $C^p$  manifold modelled on a space  $E$ , and  $F$  another Banach space. Our goal in this section is to construct the vector bundle of a manifold, which has the following desirable properties.

- The vector bundle  $W$  embeds  $X$  into itself as a submanifold.
- At each point  $p \in X$ , we attach a  $F$  space structure exclusive to each  $p$  akin to the tangent space  $T_p X$ .
- $W$  is locally isomorphic to the product space  $U \times F$ , where  $U \subseteq X$ , and
- a subset of the morphisms  $A: X \rightarrow W$  locally resemble morphisms  $U \rightarrow U \times F$ .

#### Definition 2.1 Coproduct of fibers

Suppose for each  $p$ , the set  $W_p$  is toplinearly isomorphic to  $F$  at for each  $p$  (which implies  $W_p$  is a TVS), then we call  $W_p$  an  $F$ -fiber at  $p$ . The set-theoretic coproduct of all such  $W_p$  is called a *coproduct of  $F$ -fibers modelled over  $X$* , where

$$W = \coprod_{p \in X} W_p \quad \text{comes with} \quad \pi: W \rightarrow X, \quad \pi^{-1}(p) = W_p; \quad (33)$$

with  $\pi$  being a surjection onto  $X$  called the *canonical projection* of  $W$ .

It turns out the natural way of making  $W$  a manifold would be to steal open sets from *both*  $E$  and  $F$  — in this case, sets of the form  $\hat{U} \times F$ . We sometimes write  $\widetilde{U}$  instead of  $\pi^{-1}(U)$  for brevity, and  $\tilde{p}$  in place of  $\pi^{-1}(p)$ . The next few definitions should feel familiar.

#### Definition 2.2 Local trivialisation

Let  $W$  be as in Eq. (33). A *local trivialisation* of  $W$  is a tuple  $(\widetilde{U}, \Phi)$ , such that the diagram in fig. 3 commutes, and

- $U \subseteq X$  is open in  $X$ , and for each  $p \in U$ ,
- $\Phi|_{\tilde{p}}$  is in bijection with  $W_p = F$ .

#### Definition 2.3 Compatibility between trivialisations

Let  $(\widetilde{U}, \Phi)$  and  $(\widetilde{V}, \Psi)$  be local trivialisations of  $W$ , they are called  $C^k$ -compatible if  $U \cap V = \emptyset$ , or both of the following hold:

- for each  $p \in U \cap V$  — the restriction of  $\Psi \circ \Phi^{-1}$  onto the fiber of  $p$  —  $(\Psi \circ \Phi^{-1})|_{\tilde{p}}$  is a toplinear iso-

morphism, and

- the map  $\theta: U \cap V \rightarrow L(F, F)$  as defined by Eq. (34), is a  $C^k$  morphism into the Banach space  $L(F, F)$ .

$$\theta(p) = (\Psi \circ \Phi^{-1})|_{\tilde{p}} \quad (34)$$

(equivalently, we can require  $\theta$  be a  $C^k$  morphism into the open submanifold  $\text{Laut}(F)$ ).

Note: we assume that  $0 \leq k \leq p$ .

#### Definition 2.4 Trivialisation covering

Let  $W$  be a coproduct of  $F$ -fibers over  $X$ . A  $C^k$  *trivialisation covering* of  $W$  is a collection of pairwise  $C^k$ -compatible local trivialisations  $\{(\widetilde{U}_\alpha, \Phi_\alpha)\}$  where  $\{U_\alpha\}$  is an open cover of  $X$ .

#### Definition 2.5 Vector bundle

Let  $X$  be a  $C^p$  manifold over  $E$ , and let  $F$  be a Banach space. An  *$F$ -vector bundle of rank  $k$  over  $X$*  is a coproduct of  $F$ -fibers modelled over  $X$  equipped with a **maximal  $C^k$  trivialisation covering**.

#### Remark 2.6 Maximality of trivialisation covering

One can easily verify the compatibility condition defines an equivalence relation, thus any  $C^k$ -trivialisation covering *determines* a maximal one.

#### Remark 2.7 Omissions for vector bundles

We say  $W$  is a *bundle over  $X$*  when it is unambiguous to do so.

$$\begin{array}{ccc} \widetilde{U} & \xrightarrow{\Phi} & U \times F \\ \pi \downarrow & \nearrow \text{proj}_1 & \\ U & & \end{array}$$

Figure 3: Local Trivialisation

The above definitions calls for some commentary, our end goal is to make an arbitrary rank  $C^k$  vector bundle  $W$  a  $C^k$  manifold. Open sets will still be our primary topological data. To ensure that  $W$  is as similar to  $X$  as possible, the eventual manifold structure we will put on  $W$  will **embed the structure of  $X$  into  $W$** . We are repeating (essentially) the same argument as in the submanifold case but with the roles of  $X$  and the submanifold  $S$  reversed.

Suppose we have a structure on  $W$ , then  $X = \bigcup_{p \in X} \{p\} \times 0$  is a submanifold of the  $W$  as  $E$  splits in the product space  $E \times F$ . Let us motivate a couple of the requirements above.

**Definition 2.2 (1)**  $U$  is required to be open because  $W$  inherits part of the topology, and hence the charts in  $E$  whose domain is a subset of  $U$ ,

**Definition 2.2 (2)** The second requirement implies **each  $\Phi$  is in bijection with  $\Phi(\widetilde{U}) = U \times F$ , which is open in  $E \times F$** , which will allow us to construct bijections with open subsets of the model space  $E \times F$ . Furthermore, Eq. (35) below holds for an arbitrary  $V \subseteq X$ .

$$\Phi|_{\pi^{-1}(U \cap V)} \text{ is a bijection onto } U \cap V \times F \quad (35)$$

**Definition 2.3 (1)** The overlap restricts to a top linear isomorphism on each fiber because, it allows us **to quotient out the effects of the trivialisation transitions**, by rehearsing the same 'coproduct and quotient' argument in Definitions 1.18 to 1.20.

**Definition 2.3 (2)** The requirement that the mapping Eq. (34) is a morphism, turns out to be necessary for  $\Phi_{\alpha\beta}$  to be a  $C^k$  isomorphism.

Suppose  $W$  is an  $F$ -vector bundle over  $X$  with the trivialisation covering  $\{(\widetilde{U}^\alpha, \Phi_\alpha)\}$ . For each  $\alpha$ , we can cover  $U^\alpha$  using chart domains  $(U_\beta^\alpha, \varphi_\beta^\alpha)$  in  $X$  — without loss of generality, we can assume  $U_\beta^\alpha \subseteq U^\alpha$  by restricting the chart domain and relabelling.

Similar to the construction of the induced atlas of a submanifold, given a 'piece' of the original manifold  $X$  — **instead of dropping the coordinates that correspond to  $E_2$ , we add an  $F$ -component to construct a bijection with an open subset of  $E \times F$** . This is shown in Eq. (36)

$$\tilde{\varphi}_\beta^\alpha : \widetilde{U}_\beta^\alpha \longrightarrow \hat{U}_\beta^\alpha \times F \quad \text{defined by} \quad \tilde{\varphi}_\beta^\alpha = (\varphi_\beta^\alpha \times \text{id}_F) \circ \Phi_\alpha|_{\widetilde{U}_\beta^\alpha} \quad (36)$$

### Remark 2.8 Hats and wiggles

Here,  $\widetilde{U}_\beta^\alpha$  should be interpreted as the inverse image of the open set  $U_\beta^\alpha$  through  $\pi$ . Similarly,  $\hat{U}_\beta^\alpha$  is the image of  $U_\beta^\alpha$  through  $\varphi_\beta^\alpha$ .

The collection of charts  $\mathcal{A}$ , in Eq. (37) cover  $W$  with their chart domains, and each chart is in bijection with an open subset of  $E \times F$ .

$$\mathcal{A} = \left\{ (\widetilde{U}_\beta^\alpha, \tilde{\varphi}_\beta^\alpha), (\widetilde{U}^\alpha, \Phi_\alpha) \text{ is in the trivialisation covering of } W \right\} \quad (37)$$

### Proposition 2.9 Structure of a vector bundle

Let  $X$  be a  $C^p$  manifold modelled over  $E$ . If  $W$  is a  $C^k$  vector bundle modelled on  $F$  over the manifold  $X$ , then  $W$  is a  $C^k$  manifold modelled on the product space  $E \times F$ . Furthermore:

1. The *canonical projection*  $\pi : W \rightarrow X$  is a morphism and a submersion.
2.  $X$  is  $C^k$  isomorphic to a submanifold of  $W$

*Proof.* Suppose we are given two charts in Eq. (37),  $(\widetilde{U}_{\beta_1}^{\alpha_1}, \tilde{\varphi}_{\beta_1}^{\alpha_1})$ , and  $(\widetilde{U}_{\beta_2}^{\alpha_2}, \tilde{\varphi}_{\beta_2}^{\alpha_2})$ . First, prove that  $\tilde{\varphi}_{\beta_1}^{\alpha_1}(\widetilde{U}_{\beta_1}^{\alpha_1} \cap \widetilde{U}_{\beta_2}^{\alpha_2})$  is open in  $E \times F$ :

$$\begin{aligned} \tilde{\varphi}_{\beta_1}^{\alpha_1}(\widetilde{U}_{\beta_1}^{\alpha_1} \cap \widetilde{U}_{\beta_2}^{\alpha_2}) &= \left[ (\varphi_{\beta_1}^{\alpha_1} \times \text{id}_F) \circ \Phi_{\alpha_1} \right] (\widetilde{U}_{\beta_1}^{\alpha_1} \cap \widetilde{U}_{\beta_2}^{\alpha_2}) \\ &= \left[ (\varphi_{\beta_1}^{\alpha_1} \times \text{id}_F) \circ \Phi_{\alpha_1} \right] (\pi^{-1}(U_{\beta_1}^{\alpha_1} \cap U_{\beta_2}^{\alpha_2})) \\ &= (\varphi_{\beta_1}^{\alpha_1} \times \text{id}_F)(U_{\beta_1}^{\alpha_1} \cap U_{\beta_2}^{\alpha_2}) \times F \end{aligned} \quad \text{by Eq. (35).}$$

Suppressing restrictions and computing the chart transitions in Eq. (38),

$$\tilde{\varphi}_{\beta_2}^{\alpha_2}(\tilde{\varphi}_{\beta_1}^{\alpha_1})^{-1} = (\varphi_{\beta_2}^{\alpha_2} \times \text{id}_F) \circ \Phi_{\alpha_2} \Phi_{\alpha_1}^{-1} \circ ((\varphi_{\beta_1}^{\alpha_1})^{-1} \times \text{id}_F). \quad (38)$$

which is clearly a bijection. And it is not hard to see that Eq. (38) can be factored into

$$\tilde{\varphi}_{\beta_2}^{\alpha_2}(\tilde{\varphi}_{\beta_1}^{\alpha_1})^{-1}(x, v) = \left( \varphi_{\beta_1 \beta_2}^{\alpha_1 \alpha_2}(x), [\theta \circ (\varphi_{\beta_1}^{\alpha_1})^{-1}](x)(v) \right). \quad (39)$$

for any  $x \in \varphi_{\beta_1}^{\alpha_1}(U_{\beta_1 \beta_2}^{\alpha_1 \alpha_2})$  and  $v \in F$ . **From Eq. (39), it should now be clear why we demand  $k \leq p$ .** The mapping in the second coordinate within Eq. (39) can be reduced to a composition with the evaluation map  $\mathbf{E} : \text{Laut}(F) \times F \rightarrow F$ .

$$[\theta \circ (\varphi_{\beta_1}^{\alpha_1})^{-1}](x)(v) = \mathbf{E} \circ ([\theta \circ (\varphi_{\beta_1}^{\alpha_1})^{-1}] \times \text{id}_F). \quad (40)$$

Since  $\mathbf{E}$  is continuous and bilinear, Eq. (40) and hence Eq. (38) describes a  $C^k$  mapping between open subsets of Banach spaces. It is a morphism, and reversing the roles of the two charts proves its inverse is again a morphism.

To prove  $\pi$  is a submersion, recall  $W$  is the set-theoretic disjoint union of  $F$ -fibers. Every element in  $W$  can be represented by  $(x, v) \in X \times F$ . **We will identify elements of  $W$  as elements in  $X \times F$ . However, this is not a manifold isomorphism.**

Fix  $(x, v) \in W$ , it is in the domain of some chart  $(\widetilde{U}_\beta^\alpha, \tilde{\varphi}_\beta^\alpha)$ . The  $\pi$ -image of the chart domain is  $\pi\pi^{-1}(U_\beta^\alpha) = U_\beta^\alpha$  because  $\pi$  is surjective. Using Eq. (36) and the diagram found

in fig. 3, the coordinate representation of  $\pi$  becomes

$$\begin{aligned}\pi_{(\tilde{U}_\beta^\alpha, U_\beta^\alpha)} &= \varphi_\beta^\alpha \circ \pi \circ \Phi_\alpha^{-1} \circ ((\varphi_\beta^\alpha)^{-1} \times \text{id}_F) \\ &= \varphi_\beta^\alpha \circ \text{proj}_1 \circ ((\varphi_\beta^\alpha)^{-1} \times \text{id}_F) \\ &= \text{proj}_1(\text{id}_{\tilde{U}_\beta^\alpha} \times \text{id}_F)\end{aligned}\quad (41)$$

We can differentiate both sides of Eq. (41) and if we write  $\hat{U} = \tilde{U}_\beta^\alpha$ , we obtain Eq. (42).

$$D\text{proj}_1(\text{id}_{\hat{U}} \times \text{id}_F)(x, v) = \text{proj}_1 \in L(E \times F; E) \quad \forall x \in \hat{U}, v \in F \quad (42)$$

which means  $\pi$  submersion.

Finally, the subset  $X \times 0 \subseteq W$  is easily shown to be a submanifold of  $W$ , and is isomorphic to  $X$  by dropping the  $F$  coordinate and retracing the argument we made in constructing the structure of  $W$ . ■

#### Remark 2.10 Pair of bundle charts

If  $X$  is a manifold and  $W$  a vector bundle over  $X$ , the charts realizing the representations of  $\pi$  in Eqs. (41) and (42) are called *bundle charts*.

#### Definition 2.11 Section of a vector bundle

Let  $W$  be a bundle over a manifold  $X$ . A *section* of  $W$  is a morphism  $\sigma \in \text{Mor}(X, W)$  such that the diagram in fig. 4a commutes, which is synonymous with  $\pi\sigma = \text{id}_X$ . A *local section* of  $W$  is a morphism  $\sigma : U \rightarrow W$  where  $U \subseteq X$  is viewed as a submanifold and  $\pi\sigma = \text{id}_U$ . We denote the sections of  $W$  by  $\Gamma(W)$ .

$$\Gamma(W) = \left\{ \sigma : X \rightarrow W, \sigma \text{ is a section of } W \right\} \quad (43)$$

The *zero section* of  $W$  is the section  $\sigma(p) = 0 \in W_p$  for every  $p \in X$ . If  $\sigma$  is a section of  $W$ ,  $\text{supp}(\sigma)$  refers to the *support* of  $\sigma$ , and is defined in Eq. (44).

$$\text{supp}(\sigma) = \overline{\{p \in X, \sigma(p) \neq 0\}} \quad (44)$$

#### Remark 2.12 Bundle coordinates

Let  $X$  and  $W$  be as in Def. 2.11, and suppose  $\sigma$  is a section on  $W$ . Using a pair of bundle charts,  $(U) \in X$  and  $(\tilde{U}) \in W$ , we define the *bundle coordinates* of  $\sigma$

$$\sigma_{U, \tilde{U}} = \tilde{\varphi} \circ \sigma \circ \varphi^{-1} \quad (45)$$

expanding the induced chart on  $W$  within Eq. (45) reads


$$\sigma_{U, \tilde{U}} = (\varphi \times \text{id}_F) \circ \Phi \circ \sigma \circ \varphi^{-1} \quad (46)$$

Refer to the diagram in fig. 4b. We will always use bundle charts when discussing the coordinate representa-


tion of a section, and we write

$$\sigma_U = \sigma_{U, \tilde{U}} = \hat{\sigma}$$

Sections are precisely the morphisms into  $W$  whose coordinate representation resembles that of a graph:  $\hat{\sigma} : \hat{U} \rightarrow \hat{U} \times F$  and because of this: we identify  $\hat{\sigma}(\hat{p}) = (\hat{p}, v)$  with  $v \in F$ .



(a) Section of a bundle



(b) Local coordinates of a bundle section

Figure 4: Diagrams for bundle section and its local representation

## 2.2. Tangent Bundle

#### Definition 2.13 Tangent bundle

Let  $X$  be a  $C^p$  manifold over  $E$ , the *tangent bundle* is a  $E$ -vector bundle of rank  $p-1$ , denoted by  $TX$ , and

$$TX = \coprod_{x \in X} T_x X.$$

The construction of the tangent bundle is outlined in Note 2.14.

#### Note 2.14 Construction of the Tangent Bundle

Let  $X$  be a  $C^p$  manifold with  $p \geq 1$ , so that the tangent space at every point is defined. If  $p \in (U_i, \varphi_i)$  for  $i = 1, 2$ . Then  $\varphi_{12}$  is a  $C^p$  isomorphism between  $\varphi_1(U_{12})$  and  $\varphi_2(U_{12})$ ; **whose derivative is a  $C^{p-1}$  map into  $\text{Laut}(E)$  that encodes the transformation between the concrete tangent spaces**. In the notation of Eq. (11), this means

$$x \mapsto D\varphi_{12}(x) \quad \text{is in} \quad C^{p-1}(\hat{U}_{12}, \text{Laut}(E)).$$

In fact, the tangent bundle  $TX \stackrel{\Delta}{=} \coprod_{p \in X} T_p X$  is a  $C^{p-1}$  vector bundle (modelled on  $E$ ) over  $X$ . If  $(U, \varphi)$  is a

chart in  $X$ , it induces a local trivialisation on  $TX$  by taking each tangent vector  $v \in T_p X$  to its concrete representation  $(p, \hat{v}) \in X \times E$ .

$$\Phi: \widetilde{U} \rightarrow U \times E \quad \text{and} \quad \Phi(v) = (p, \hat{v}) \quad (47)$$

where  $(U, \varphi, p, \hat{v})$  is a concrete representation of  $v \in T_p X$ .

Similarly, we have the cotangent bundle which is modelled on toplinear dual of the tangent spaces of  $X$ .

### Definition 2.15 Cotangent bundle

Let  $X$  be a  $C^p$  manifold over  $E$ , the *cotangent bundle* is a  $E$ -vector bundle of rank  $p-1$ , denoted by  $T^* X$ , and

$$T^* X = \bigsqcup_{x \in X} T_x X^*,$$

where  $T_x X$  is toplinearly isomorphic to  $X$ , and  $T_x X^*$  its toplinear dual.

## 2.3. Notes and References

### Category Theory

Let  $X$  and  $E_i$  be Banach spaces for  $i = \underline{2}$ . We discussed the difference in the role that a toplinear mapping  $f \in L(E_1, E_2)$  plays in pushing points from  $E_1 \rightarrow E_2$ , and the role it plays from pushing *incoming maps* with source  $X$  from  $L(X, E_1) \rightarrow L(X, E_2)$  by composing the incoming map with itself.

$$f: E_1 \rightarrow E_2 \quad \text{and} \quad f_c: L(X, E_1) \rightarrow L(X, E_2) \quad (48)$$

where the  $c$  within  $f_c$  stands for composition. This is summarized in fig. 5

$$\begin{array}{ccc} X & \xrightarrow{\lambda} & E_2 \\ & \searrow \lambda_c(f) & \downarrow f \\ & & E_2 \end{array}$$

Figure 5: Functorality through post-composition

The two maps  $f$  and  $f_c$  often thought of as the same, and we can identify  $f$  with two separate actions. Here, it is not so obvious why we need the concept of functorality, a closer look at the **different roles that the same mathematical object can play** will surely motivate the above discussion.

Let  $E_i$  be Banach spaces,  $f \in C^p(E_1, E_2)$  and  $\lambda \in L(E_2, E_3)$ .

- The derivative of  $f$  is a continuous map  $Df: E_1 \rightarrow L(E_1, E_2)$ .

- The derivative of  $\lambda$  (now 'identified with  $\lambda \in C^p(E_2, E_3)$ ), is continuous constant map from  $E_2$  to  $L(E_2, E_3)$

$$D(\lambda \circ f) = \lambda \circ Df(x) \quad (49)$$

but what if we have a multi-linear map whose destination is  $E_1$ , and what about symmetric/alternating multi-linear maps, continuous maps,  $C^p$  morphisms? Should we let  $f$  take on all of those roles as well? Should we identify  $f$  with its adjoint as well? This is the first of the many problems.

The problem becomes even clearer when we look at maps between  $F$ -fibers. Fix a manifold  $X$  and  $F_i$ -bundles  $W^i$  over  $X$  for  $i = \underline{2}$ . Suppose  $A: X \rightarrow W^1$  is a section on  $W^1$ , and  $\lambda \in L(F_1, F_2)$ .

At each point  $p \in X$ , our linear map  $\lambda$  can be identified with the linear map that acts between the fibers.

$$\lambda_p: W_p^1 \rightarrow W_p^2$$

which is toplinear hence a morphism. Our main problem is concerned with the conditions under which the composition  $\lambda A$  — as defined in Eq. (50) — is a morphism.

$$\lambda A: X \rightarrow W^2 \quad \text{and} \quad (\lambda A)(p) = \lambda_p(A_p) \in W_p^2 \quad (50)$$

Under what conditions can a morphism take on additional roles? The mapping  $\lambda_p$  on each tangent space is a  $C^p$  morphism in the manifold sense, and the morphisms that preserve the  $C^p$  smoothness of sections are called VB morphisms. Which we will define after some more category theory.

For the remainder of this section, let  $C_1$  and  $C_2$  be categories. We denote the objects of  $C_1$  by  $E_i$ , and the objects of  $C_2$  by  $F_i$ .

### Definition 2.16 Functor

A correspondence  $\theta$  between  $C_1$  and  $C_2$  is called a *functor* — which we denote by  $\theta: C_1 \Rightarrow C_2$  — if all of the following rules satisfied.

**Ob1**  $\theta$  maps objects in  $C_1$  to objects in  $C_2$ . We write

$$\forall E \in Ob(C_1), \quad E^\theta = \theta(E) \in Ob(C_2) \quad (51)$$

**Mor1**  $\theta$  associates morphisms in  $C_1$  to morphisms in  $C_2$  that respects Ob1.

$$\forall f \in Mor_{C_1}(E_1, E_2), \quad \theta(f) \in Mor_{C_2}(E_1^\theta, E_2^\theta) \quad (52)$$

**Mor2** Identity is associated with identity:  $\theta(id_E) = \theta(id_{\theta(E)})$ .

**Mor3** Commutes with inversion:  $\theta(f^{-1}) = \theta(f)^{-1}$  if the inverse of  $f$  exists.

**Mor4** Functorial:  $\theta(g \circ f) = \theta(g) \circ \theta(f)$ .

### Note 2.17 The $\text{Hom}_X$ functor

We continue our discussion from fig. 5. Recall  $\text{Ban}_{\mathbb{R}}$  is the category of Banach spaces over  $\mathbb{R}$ , and we will refer to toplinear morphisms as morphisms for brevity. If  $X$  is an object in  $\text{Ban}_{\mathbb{R}}$ , the  $\text{Hom}_X$  functor is a covariant functor between  $\text{Ban}_{\mathbb{R}}$  and  $L(X, \cdot)$  — the space of toplinear mappings with source  $X$  such that

1. to each  $E_i \in \text{Ob}(\text{Ban}_{\mathbb{R}})$   $\text{Hom}_X(E_i) = L(X, E_i)$  — **the space of incoming morphisms with source  $X$** , and
2. to each morphism  $f \in L(E_1, E_2)$  another morphism between  $L(X, E_1)$  and  $L(X, E_2)$  — denoted by  $(\text{Hom}_X)f$ .
3. The functor  $\text{Hom}_X$  converts **outgoing morphisms from  $E_1$  to the redirection morphism of incoming morphisms with source  $X$** .

Notice this is precisely what the diagram in fig. 5 describes.

### Proposition 2.18 $\text{Hom}_X$ functor is a functor

The  $\text{Hom}_X$  functor as defined in Note 2.17 is a covariant functor.

*Proof.* Postponed. ■

### Note 2.19 The tangent space functor

Let  $X$  be a  $C^1$  manifold, we call the tuple  $(p, X)$  for  $p \in X$  the *centering of  $X$  centered at  $p$* . The category of pointed manifolds, denoted by  $\text{Man}_*$ . Its objects consist of all centerings across  $C^1$  manifolds, and the morphisms in  $\text{Man}_*$  are called *pointed morphisms*.

If  $(q, Y)$  is another object in  $\text{Man}_*$ , a pointed morphism between  $(p, X)$  and  $(q, Y)$  is a tuple  $(p, f)$ ; where  $f$  is a manifold morphism between  $X$  and  $Y$  and  $f(p) = q$ . We sometimes write  $f_p = (p, f)$  when it is clear.

The *tangent space functor*, denoted by  $T : \text{Man}_* \Rightarrow \text{Ban}_{\mathbb{R}}$  is a covariant functor where

- we define  $T(p, X) = T_p X$  that takes a pointed  $C^1$  manifold to its tangent space, and
- to each pointed  $C^1$  morphism  $f_p \in \text{Mor}_{\text{Man}_*}((p, X), (q, Y))$  we associate with the

toplinear mapping

$$Tf_p = df(p) : T_p X \rightarrow T_q Y \quad (53)$$

### Proposition 2.20 Tangent space functor is a functor

The tangent space functor as defined in Note 2.19 is a covariant functor.

*Proof.* Postponed. ■

We leave the verification that  $T$  satisfies the properties in Def. 2.16 as an exercise. Dual to the concept of a functor is that of the cofunctor, which — for our purposes — captures the idea of the toplinear adjoint.

### Definition 2.21 Cofunctor

Let  $C_1$  and  $C_2$  be categories. A correspondence  $\eta : C_1 \leftrightarrows C_2$  is called a *cofunctor* (or a contravariant functor) if all of the following rules are satisfied.

**Ob**  $\eta$  maps objects in  $C_1$  to objects in  $C_2$ . We write  $E^\lambda = \lambda(E) \in \text{Ob}(C_2)$  for every  $E \in \text{Ob}(C_1)$ .

**Mor1**  $\eta$  associates morphisms in  $C_1$  to morphisms in  $C_2$  that respects  $\text{Ob}$ .

$$\forall f \in \text{Mor}_{C_1}(E_1, E_2), \quad \eta(f) \in \text{Mor}_{C_2}(E_2^\eta, E_1^\eta) \quad (54)$$

**Mor2** Identity is associated with identity:  $\eta(\text{id}_E) = \eta(\text{id}_{\eta(E)})$ .

**Mor3** Commutes with inversion:  $\eta(f^{-1}) = \eta(f)^{-1}$  if the inverse of  $f$  exists.

**Mor4** Cofunctoral:  $\eta(g \circ f) = \eta(f) \circ \eta(g)$ .

### Remark 2.22 Cofunctors are opposite to functors

The cofunctor  $\eta$  reverses the arrows a morphism  $f$ . Refer to fig. 6 for a comparison between Eq. (54) and Eq. (52).

### Note 2.23 The $\text{Hom}^X$ cofunctor

Let  $X \in \text{Ob}(\text{Ban}_{\mathbb{R}})$ , it defines a cofunctor from  $\text{Ban}_{\mathbb{R}} \leftrightarrows L(\cdot, X)$  where  $L(\cdot, X)$  is the space of toplinear mappings whose destination is  $X$ .

1. to each  $E_i \in \text{Ob}(\text{Ban}_{\mathbb{R}})$   $\text{Hom}^X(E_i) = L(E_i, X)$  — **the space of outgoing morphisms with destination  $X$** , and
2. to each morphism  $f \in L(E_1, E_2)$  another morphism between  $L(E_1, X)$  and  $L(E_2, X)$  — denoted by  $(\text{Hom}^X)f$ .
3. The functor  $\text{Hom}^X$  converts **outgoing morphisms from  $E_1$  to the precomposition morphism with destination  $X$** .

**phism which acts on morphisms with destination  $X$ .**

$$\begin{array}{ccccc}
 \eta(E_1) & \xleftarrow{\eta} & E_1 & \xrightarrow{\theta} & \theta(E_1) \\
 \uparrow \eta(f) & & \downarrow f & & \downarrow \theta(f) \\
 \eta(E_2) & \xleftarrow{\eta} & E_2 & \xrightarrow{\theta} & \theta(E_2)
 \end{array}$$

Figure 6: Functor  $\theta$  vs. cofunctor  $\eta$  comparison

### 3. Coordinates

#### 3.1. Introduction

In the previous chapters, a chart  $(U, \varphi)$  was often equated with its domain. We will now express a concrete tangent vector as  $(\hat{p}, \hat{v})$ , omitting any reference to the chart or its domain.

Let  $X$  be a manifold and  $F$  a Banach space. Consider a morphism  $f \in \text{Mor}(X, F)$  and fix a point  $p \in X$ , and write  $q = f(p)$ . By adopting the canonical interpretation  $\bar{w}$  for a tangent vector  $w \in T_q F$  (as discussed in Rmk 1.23), we

- reinterpret the differential at  $p$   $df_p$  as a linear map from  $T_p X$  to  $F$ ,
- always use the standard chart  $(\text{id}_F, F)$  so that  $\hat{f} = f_{U,F}$ .

In this context, morphisms into  $\mathbb{R}$  almost serve as test functions in the framework of distribution theory. This requires a definition.

#### Definition 3.1 Function on $X$

Let  $X$  be a manifold of class  $C^p$  over  $\mathbb{R}^n$  for  $n, p \geq 1$ . A *function* on  $X$  is a morphism  $f : X \rightarrow \mathbb{R}$ , where  $\mathbb{R}$  should be interpreted as a manifold. We denote the commutative ring of functions on  $X$  by  $C^p(X, \mathbb{R})$  or  $C^p(X)$ . If  $U$  is an open subset of  $X$ , its functions are denoted by  $C^p(U, \mathbb{R})$  or  $C^p(U)$ .

#### 3.2. Exterior Derivative

Let  $X$  be a manifold, and  $f \in C^p(X)$ . If  $\gamma : (-\delta, +\delta) \rightarrow X$  is a curve starting at  $x_0 \in X$  with velocity  $v$ , the composition  $f \circ \gamma$  is a morphism. Let us write

$$F : (-\delta, +\delta) \rightarrow \mathbb{R}, \quad F(\varepsilon) = f \circ \gamma(\varepsilon) - f \circ \gamma(0) \quad (55)$$

Suppose we wish to measure the rate at which  $f$  moves in the direction of  $v$ , then we can simply take the derivative

of Eq. (55). We define the *exterior derivative of  $f$  at  $x_0$* , denoted by  $df(x_0) : T_{x_0} X \rightarrow \mathbb{R}$  by Eq. (56)

$$df(x_0)(v) = DF(0)(\bar{1}) \quad \text{where } F = f \circ \gamma \quad (56)$$

for any curve starting at  $x_0$  with velocity  $v$ .

Let  $E$  be a Banach space, and suppose  $\omega$  is a  $k$ -form on  $E$ , and  $x_0 \in E$  with  $k+1$  tangent vectors  $v_{\underline{k+1}}$ . The parallelopiped defined by the  $k+1$  vectors is

$$P_{x_0}(v_{\underline{k+1}}) = \left\{ x_0 + \sum_{i=\underline{k+1}} t_i v_i, 0 \leq t_i \leq 1 \forall i = \underline{k+1} \right\}$$

As with Eq. (55), we can integrate over the boundary defined by  $P_{x_0}(v_{\underline{k+1}})$ , and obtain a new function. We can shrink each  $v_i$  by  $\varepsilon_i$ , and we define

$$P_{x_0}^{\varepsilon_{\underline{k+1}}}(v_{\underline{k+1}}) = \left\{ x_0 + \sum_{i=\underline{k+1}} t_i v_i, 0 \leq t_i \leq \varepsilon_i \forall i = \underline{k+1} \right\} \quad (57)$$

(Note: Perhaps after shrinking the domain of  $F$ , here we should replace everything by their coordinate representations).

$$F : (-\delta, +\delta)^{k+1} \rightarrow \mathbb{R}, \quad F(\varepsilon_{\underline{k+1}}) = \int_{\partial P_{x_0}^{\varepsilon_{\underline{k+1}}}(v_{\underline{k+1}})} \omega \quad (58)$$

We define the exterior derivative of a  $k$ -form by the map  $DF(0)(1^{(k+1)})$ .

#### 3.3. Derivations

For the rest of this chapter, assume all manifolds to be  $C^\infty$  manifolds over  $\mathbb{R}^n$ .

#### Definition 3.2 Algebra of germs

Let  $X$  be a manifold, and  $p \in X$ . A *germ at  $p$*  is an equivalence class of smooth functions, under the relation  $f \sim g$  iff  $f = g$  on some neighbourhood of  $p$ . The *algebra of germs at  $p$*  is denoted by  $C_p^\infty(X)$ , under pointwise multiplication.

#### Definition 3.3 Derivation at a point

A *derivation* at a point  $p$  is a linear functional  $v \in C_p^\infty(X)^*$  on the space of germs, that satisfies the product rule:

$$v(m(f_{\underline{k}})) = \sum_{i=\underline{k}} m(f_{i-1}(p), v f_{i-1}, f_{i+\underline{k}-i}),$$

where  $m$  denotes pointwise multiplication. The set of derivations at  $p$  is denoted by  $\mathcal{D}_p X$ .

We turn to the case in  $\mathbb{R}^n$ . Let  $\hat{U}$  be an open subset of  $\mathbb{R}^n$ , and  $\hat{p} \in \hat{U}$ . To every smooth function  $\hat{f} \in C^\infty(\hat{U}, \mathbb{R})$ , and  $\hat{v} \in T_{\hat{p}} \hat{U} = \mathbb{R}^n$ , we define

$$\langle \hat{v}, \hat{f} \rangle_{T_{\hat{p}} \hat{U}} = D\hat{f}(\hat{p})(\hat{v}).$$

We remark that the right member can be computed using  $\langle \hat{v}, \text{grad } \hat{f}(\hat{p}) \rangle_{\mathbb{R}^n}$ . Furthermore,  $\hat{f} = \hat{g}$  on some neighbourhood about  $\hat{p}$ , iff  $\langle \hat{v}, \hat{f} - \hat{g} \rangle_{T_{\hat{p}}\hat{U}} = 0$ ; and the product rule tells us that  $\langle \hat{v}, \cdot \rangle_{T_{\hat{p}}\hat{U}}$  is a derivation. On the other hand, the components of  $\hat{v}$  can be computed by deriving the coordinate functions:  $\langle \hat{v}, e_i \rangle_{T_{\hat{p}}\hat{U}} = \hat{v}^i$ .

#### Proposition 3.4

Let  $X$  be a manifold and  $p \in X$ . To every tangent vector  $v \in T_p X$ , and  $f \in C_p^\infty(X)$ ,

$$\langle v, f \rangle_{T_p X} = Df_{U, \mathbb{R}}(\hat{p})(\hat{v})$$

Let  $E$  and  $F$  be Banach spaces and  $U \subseteq E$ , suppose  $f$  is a morphism from  $U$  to  $F$ . If  $p$  is a point in  $U$ ,  $Df(p)$  is of course a linear map from  $E$  to  $F$ ; this suggests a natural pairing  $\hat{\mathcal{D}}$  of  $f$  with and  $(p, v) \in U \times E$  as shown in Eq. (59).

$$\hat{\mathcal{D}} : (U \times E) \times C^k(U, F) \longrightarrow F: \quad ((p, v), f) \mapsto Df(p)(v) \in F \quad (59)$$

Suppose  $F = \mathbb{R}$  and denote pointwise multiplication on  $\mathbb{R}$  by  $m$ . The above pairing trivially satisfies the product rule displayed in Eq. (60).

$$Dm(f_{\underline{k}})(p)(v) = \sum_{i=k} m(f_{i-1}(p), Df_i(p)(v), f_{i+k-i}(p)) \quad (60)$$

where  $f_{\underline{k}} \in C^p(U, \mathbb{R})$ . Next, if  $f$  is a function (from a manifold  $X$ ) defined on an open neighbourhood  $U$  of  $p$ . If  $v \in T_p X$ , the commentary in the introduction suggests a 'duality pairing' between  $f$  and  $(p, v)$  in the form of Eq. (61).

$$\mathcal{D} : (U \times E) \times C^p(U, F) \longrightarrow F \quad \text{and} \quad \mathcal{D}((p, v), f) = df_p(v) \quad (61)$$

By definition of the differential  $df_p$ , the right hand side of Eq. (61) is representation independent, hence

$$\mathcal{D}((p, v), f) = D\hat{f}(\hat{p})(\hat{v}), \quad \text{where the right member is an ordinary derivative of } \hat{f} \text{ at } \hat{p}, \quad (62)$$

for any representation  $(\hat{p}, \hat{v})$ ,  $\hat{f}$ . We also see that  $\mathcal{D}((p, v), f) = \hat{\mathcal{D}}((\hat{p}, \hat{v}), \hat{f})$ , which shows functions defined on  $U$  are dual to  $T_p X$  for each  $p \in U$ . We will make this notion precise when we introduce covectors.

#### Definition 3.5 Derivation at $p$

A derivation at  $p$  is a linear functional  $v$  on  $C^p(U, \mathbb{R})$ , where  $U$  is any neighbourhood of  $p$ ; such that for  $f_{\underline{k}} \in C^p(U)$ , Eq. (63) holds.

$$v(m(f_{\underline{k}})) = \sum_{i=k} m(f_{i-1}(x), v(f_i), f_{i+k-i}(x)) \quad (63)$$

We will denote the space of derivations at  $p$  by  $\mathcal{D}_p(X)$ , and if  $v \in \mathcal{D}_p(X)$ , we say  $v$  derives  $f$  for any function  $f$  defined about  $p$ .

We have shown every tangent vector is a derivation, since the product rule descends from Eq. (60) and its computation in coordinates in Eq. (62). If  $X$  is finite-dimensional, Prop. 3.6 shows derivations at a point  $p \in X$  are uniquely represented by a tangent vector.

#### Proposition 3.6

Let  $p$  be a point on a manifold  $X$ , then its tangent space is isomorphic to the vector space of derivations. If  $(\hat{p}, \hat{v})$  is a concrete tangent vector, its derivation of  $f$  computed using Eq. (62).

*Proof.* Postponed. ■

### 3.4. Boundary

#### Definition 3.7 Subsets of the upper half-plane

Let  $n \geq 1$ , the *upper half plane* of  $\mathbb{R}^n$  is the subset  $\mathbb{H}^n = [x^n \geq 0]$ . Its topological interior (resp. boundary) is denoted by  $\text{Int } \mathbb{H}^n = [x^n > 0]$  (resp.  $\partial \mathbb{H}^n = [x^n = 0]$ ).

We wish to define a notion of the boundary for a manifold. Instead of modelling  $X$  on open subsets of  $\mathbb{R}^n$ , we consider open subsets of both  $\mathbb{R}^n$  and  $\mathbb{H}^n$ . To wit, we start by generalizing the notion of  $C^p$  smoothness between **arbitrary subsets of  $\mathbb{R}^n$** . Let  $S \subseteq \mathbb{R}^n$  be a subset of  $\mathbb{R}^n$ , recall that:

A function  $f : S \rightarrow \mathbb{R}^m$  is said to be continuous on  $S$ , if  $f^{-1}(U)$  is relatively open in  $S$  for every  $U \subseteq \mathbb{R}^m$ .

The take-away intuition for this is that manifolds with boundary are supposedly used to model geometric objects that are suddenly 'cut off'. In the case of the upper half-plane, this manifests in a sudden stop in the  $n$ -th coordinate. The morphisms we seek are the ones which are ordinary morphisms but whose domains 'cut off'.

Let  $F : S \rightarrow S'$  be a mapping between arbitrary subsets of  $\mathbb{R}^n$  and  $\mathbb{R}^m$ . It is a  $C^p$  morphism, whenever each point  $p \in S$  admits a  $C^p$  extension to a neighbourhood containing  $p$ .

With this, a  $C^p$  isomorphism between subsets of  $\mathbb{R}^n$  is a bijective  $C^p$  morphism whose inverse is also a  $C^p$  morphism.

#### Definition 3.8 Boundary chart

Let  $X$  be a non-empty subset. A *boundary chart* on  $X$  modelled on  $\mathbb{R}^n$  is a tuple  $(U, \varphi)$ , such that  $U \subseteq X$ ,  $\varphi(U) = \hat{U}$  is an open subset of  $\mathbb{H}^n$ , and  $\varphi$  is a bijection onto  $\hat{U}$ .

To distinguish between this new definition of the previous one, the word *chart* will always refer to the charts defined in Chapter 1. If we wish to be precise, we will use the word *interior chart*.

**Definition 3.9 Compatibility between boundary and interior charts**

Let  $(U, \varphi)$  and  $(V, \psi)$  be boundary or interior charts of  $X$ , modelled on  $\mathbb{R}^n$  or  $\mathbb{H}^n$ . They are called  $C^p$ -compatible (for  $p \geq 0$ ) if  $U \cap V = \emptyset$ , or both of the following hold:

- $\varphi(U \cap V)$  and  $\psi(U \cap V)$  are *both* open subsets of  $\mathbb{R}^n$  or  $\mathbb{H}^n$ .
- the *transition map*  $\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V)$  is a  $C^p$  isomorphism between open subsets of  $\mathbb{R}^n$  or  $\mathbb{H}^n$ .

**Definition 3.10 Boundary atlas, structure with boundary**

Let  $X$  be a non-empty set and  $p \geq 0$ . A  $C^p$  *boundary atlas* on  $X$  modelled on  $\mathbb{R}^n$  is a pairwise  $C^p$ -compatible collection of charts  $\{(U_\alpha, \varphi_\alpha)\}$  whose union over the domains cover  $X$ .

If  $\mathcal{A}$  is a boundary atlas of  $X$ , the maximal boundary atlas containing  $\mathcal{A}$  is called the  $C^p$  *boundary structure determined by*  $\mathcal{A}$ . A  $C^p$  *manifold with boundary* modelled on  $\mathbb{R}^n$  is a non-empty set  $X$  with a  $C^p$  boundary structure modelled on  $\mathbb{R}^n$ .

We also have the following terminology for points in a manifold with boundary  $X$ . Let  $p \in X$ ,

- $p$  is called a *boundary point* whenever it is in the domain of a boundary chart  $(U, \varphi)$ , and  $\varphi(p) \in \partial\mathbb{H}^n$ ,
- $p$  is called an *interior point* otherwise.

**Remark 3.11**

It can be shown (using deRham cohomology, see Chapter 16-17 [Lee13]) that a point on a manifold cannot be a boundary point and an interior point at the same time.

## References

- [Lan12] Serge Lang. *Fundamentals of Differential Geometry*. Springer Science & Business Media, Dec. 2012.
- [Lee13] John M. Lee. *Introduction to Smooth Manifolds*. 2nd ed. Springer Science & Business Media, Mar. 2013.